



Section 18
Emergency
Exemption

FOR
DISTRIBUTION
AND USE ONLY IN
WISCONSIN

4F Herbicide

FMC Corporation, Agricultural Products Group

For Agricultural or Commercial Use Only

EPA Reg. No. 279-3220

EPA File Symbols: 07-WI-02

EMERGENCY CALLS: 800-331-3148

Contains 4 pounds of active ingredient (39.6%) per gallon.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS OF THIS PRODUCT LABEL MUST BE FOLLOWED.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

This exemption is effective from June 20, 2007 through December 15, 2007.

Crop	Weeds Controlled		Rate of Application
Strawberry	Common groundsel Common lambsquarters Corn spurry Ladysthumb Mayweed	Redroot pigweed Pineapple weed Prostrate knotweed Wild buckwheat	4 to 8 ounces per acre 0.125 to 0.25 lb ai per acre

DIRECTIONS FOR USE:

Apply in 20 to 40 gallons per acre of water by ground application only.

Apply 4 to 8 fluid ounces of formulated product (0.125 to 0.25 lbs ai) per acre per application. A maximum of 12 fluid ounces (0.375 lbs ai) can be applied per acre per season. Apply either a single application of no more than 8 fluid ounces (0.25 lbs ai) of Spartan® 4F Herbicide per acre; or apply two split applications at renovation and dormancy applying no more than a total of 12 fluid ounces (0.375 lbs ai).

Sulfentrazone may be applied during two distinct time periods:

- From June 20, 2007 through July 20, 2007, after harvest and immediately after mowing, and
- From October 15, 2007 through December 15, 2007 during dormancy.

It is a violation of federal law to harm any endangered species. Growers/applicators uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection's Biologist, at 608-224-4538, if assistance is needed to identify or protect a species or their habitat.

PRECAUTION:

Applications to strawberry plants with newly emerged growth are not recommended due to leaf burning and possible stand loss.

After spraying Spartan 4F and before using sprayer equipment for any other application, the sprayer must be thoroughly cleaned using the procedure on the EPA label. Failure to remove even small amounts of Spartan 4F from application equipment may result in injury to subsequent sprayed crops.

GENERAL INFORMATION:

Spartan 4F is a selective soil-applied herbicide for the control of certain broadleaf weeds, grasses and sedges. Spartan 4F is a suspension concentrate formulation containing 4 pounds of the active ingredient, sulfentrazone, per gallon.

The mode of action of Spartan 4F involves uptake by weed roots and shoots. Preemergence applications of Spartan 4F require rainfall or irrigation to activate the herbicide. The amount of rainfall or irrigation required for activation following application depends on existing soil moisture, organic matter content and soil texture. If adequate moisture of 0.5 to 1 inch is not received within 7 to 10 days after the application of Spartan 4F, a shallow cultivation may be needed to obtain desired weed control.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Spartan and **FMC** -FMC Trademarks
©2007 FMC Corporation

ROTATIONAL GUIDELINES:

Long rotational periods are recommended for some crops following applications of Spartan 4F. Refer to the EPA label for specific guidelines.

RESTRICTIONS:

Do not exceed 8 ounces, 0.25 lb ai, per acre per application or a maximum of 12 ounces, 0.375 lb ai, per acre per season.

Do not apply within 105 days of harvest.

Observe the restricted entry interval of 12 hours. Follow PPE guidelines on the EPA label.

Application by air or through any type of irrigation system (ie chemigation) is prohibited.

Do not use flood irrigation to incorporate this product.

Do not apply to soils classified as sand with less than 1 percent organic matter.

Do not apply to saturated soils. If soil is approaching saturation, avoid application when heavy rainfall is predicted to occur within 24 hours following planned application

This product is toxic to marine and estuarine invertebrates. Do not apply directly to water or wetlands, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

ENDANGERED SPECIES PROTECTION REQUIREMENT

Endangered species are present throughout the state of Wisconsin. It is a violation of federal law to harm an endangered species. WDATCP operates an Endangered Species program providing consultation to growers throughout the state. If a grower is uncertain of the presence of a protected species, they should contact the WDATCP Endangered Species biologist at 608-224-4538, for assistance.

To protect listed threatened or endangered (T/E) plant species, one of the following options must be followed:

1. Apply only when there is sustained wind away from native plant communities,
2. Leave a 25 foot untreated buffer between treatment area and native plant communities, or
3. Use low-pressure nozzles according to manufacturer's specifications that produce only coarse or very coarse droplets (no less than 350 micron).

SPECIAL PRECAUTIONS:

Proper Handling Instructions: This product may not be mixed or loaded within 50 feet of any wells, including abandoned wells and drainage wells, sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing and loading areas. See guidelines on the EPA label pertaining to impervious pads and diked mixing and loading areas. The product must be used in a manner that will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Groundwater Advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. **Do not use on coarse soils classified as sand, which have less than 1 percent organic matter.**

Surface Water Advisory: Sulfentrazone can contaminate surface water through spray drift. Under some conditions sulfentrazone may also have a high potential for runoff into surface water, primarily via dissolution in runoff water, for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

CONDITIONS OF USE:

FMC Corporation intends that this Section 18 label be distributed only to end users and/or growers and/or applicators acting on behalf of growers who agree that FMC would not be held responsible for failure to perform and/or any crop damage from use of Spartan Herbicide in pre-plant, post-transplant, renovation or maintenance treatments on strawberry. If such terms and conditions are unacceptable, FMC requests the return of Spartan Herbicide at once unopened. FMC considers the user to have accepted such terms and conditions upon use of this product.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED SPARTAN 4F AND THE EMERGENCY EXEMPTION SUPPLEMENTAL LABEL MUST BE FOLLOWED.

THE WISCONSIN DEPARTMENT OF AGRICULTURE MUST BE INFORMED OF ANY ADVERSE EFFECTS, WHICH MAY RESULT FROM THE USE OF THIS PRODUCT.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103
Spartan4F_18_WI_straw_03-12-2007

Spartan and  -FMC Trademarks
©2007 FMC Corporation